

173	-	1	2	3	Muscle weakness
174	-	1	2	3	Lack of Stamina
175	-	1	2	3	Drowsiness after eating
176	-	1	2	3	Muscular soreness
177	-	1	2	3	Rapid heart beat
178	-	1	2	3	Hyper-irritable
179	-	1	2	3	Feeling of a band around your head
180	-	1	2	3	Melancholia (feeling of sadness)
181	-	1	2	3	Swelling of ankles
182	-	1	2	3	Diminished urination
183	-	1	2	3	Tendency to consume sweets or carbohydrates
184	-	1	2	3	Muscle spasms
185	-	1	2	3	Blurred vision
186	-	1	2	3	Loss of muscular control
187	-	1	2	3	Numbness
188	-	1	2	3	Night sweats
189	-	1	2	3	Rapid digestion
190	-	1	2	3	Sensitivity to noise
191	-	1	2	3	Redness of palms of hands and bottom of feet
192	-	1	2	3	Visible veins on chest and abdomen
193	-	1	2	3	Hemorrhoids
194	-	1	2	3	Apprehension (feeling that something bad is going to happen)
195	-	1	2	3	Nervousness causing loss of appetite
196	-	1	2	3	Nervousness with indigestion
197	-	1	2	3	Gastritis
198	-	1	2	3	Forgetfulness
199	-	1	2	3	Thinning hair

200	–	1	2	3	Very easily fatigued
201	–	1	2	3	Premenstrual tension
202	–	1	2	3	Painful menses
203	–	1	2	3	Depressed feelings before menstruation
204	–	1	2	3	Menstruation excessive and prolonged
205	–	1	2	3	Painful breasts
206	–	1	2	3	Menstruate too frequently
207	–	1	2	3	Vaginal discharge
208	–	1	2	3	Hysterectomy/ovaries removed
209	–	1	2	3	Menopausal hot flashes
210	–	1	2	3	Menses scanty or missed
211	–	1	2	3	Acne, worse at menses
212	–	1	2	3	Depression of long standing

213	—	1	2	3	Prostate trouble
214	—	1	2	3	Urination difficult or dribbling
215	—	1	2	3	Night urination frequent
216	—	1	2	3	Depression
217	—	1	2	3	Pain on inside of legs or heels
218	—	1	2	3	Feeling of incomplete bowel evacuation
219	—	1	2	3	Lack of energy
220	—	1	2	3	Migrating aches and pains
221	—	1	2	3	Tire too easily
222	—	1	2	3	Avoids activity
223	—	1	2	3	Leg nervousness at night
224	—	1	2	3	Diminished sex drive

TO THE PATIENT: Please list below the five main physical complaints you have in order of their importance.

1. _____
2. _____
3. _____
4. _____
5. _____

Postural Blood Pressure: Recumbent _____ Standing _____ Pulse _____

Hema-Combistix Urine readings: pH _____ Albumin per cent _____ Glucose per cent _____

Occult Blood _____ pH of Saliva _____ pH of Stool specimen _____ Weight _____

Hemoglobin _____ Blood Clotting Time _____

This test was developed by Dr. Broda Barnes, M.D. and is a measurement of the underarm temperature to determine hypo and hyperthyroid states. The test is conducted by the patient in the a.m. before leaving bed - with the temperature being taken for 10 minutes. The test is invalidated if the patient expends any energy prior to taking the test - getting up for any reason, shaking down the thermometer, etc. It is important that the test be conducted for exactly 10 minutes, making the prior positioning of both the thermometer and a clock important.

Any two days during the month

The 2nd and 3rd day of flow OR any 5 days in a row.

Any 2 days during the month.

You can do the following test at home to see if you may have a functional low thyroid. Use an oral thermometer or a digital one. When you use a digital one, place the probe under your arm for 5 minutes then turn your machine on; continue on for an additional 5 minutes. When using a regular one, shake down the night before.

[illegible]